



Review and Approval 5/14/2018, Effective 8/1/18, replaces all prior versions **PROTOCOL 17A: Destination Determination – Adult General Medical Patients**






Review and Approval 5/14/2018, Effective 8/1/18, replaces all prior versions **PROTOCOL 17A: Destination Determination – Adult Cardiology Patients**






Review and Approval 5/14/2018, Effective 8/1/18, replaces all prior versions **PROTOCOL 17A: Destination Determination – Adult Stroke Patients**











Review and Approval 5/14/2018, Effective 8/1/18, replaces all prior versions **PROTOCOL 17A: Destination Determination– Pediatric General Medical Patients**






Review and Approval 5/14/2018, Effective 8/1/18, replaces all prior versions **PROTOCOL 17A: Destination Determination – Pediatric Trauma Patients**





Review and Approval 5/14/2018, Effective 8/1/18, replaces all prior versions





| _ | | Neonatal Priority Determination | |
|---|-----------------------|--|--|
| | | | |
| • | Priority I - Unstable | | |
| | | Cardiac or respiratory arrest | |
| | | Less than 35 weeks gestation at time of birth(estimated) | |
| | | APGAR ≤ 5 at 5 minutes | |
| | | SpO2 less than 90% on oxygen | |
| | | Diagnosed genetic disorders | |
| | | | |
| | | | |
| | | | |
| • | Prio | Priority II - Stable | |
| | | 35 weeks or later gestation at time of birth | |
| | | APGAR > 5 at 5 minutes | |
| | | No immediate life threat identified | |
| | | | |